


DTIC FILE COPY

UNCLASSIFIED UNLIMITED

②

ARE TM (UDT) 89114



ARE TM (UDT) 89114

ACCN. No. 79462

AUGUST 1989

COPY No. 16

AD-A220 053

## A NOTE ON THE GAUSSIAN INTEGRATION FORMULA

by  
C RICHARDSON

ACCN. No. 79462

DTIC  
JAN 0 1990  
E

ADMIRALTY RESEARCH ESTABLISHMENT  
Procurement Executive, Ministry of Defence,  
Southwell, Portland, Dorset.

UNCLASSIFIED UNLIMITED

04 04 103

### CONDITIONS OF RELEASE

- A**

## CONDITIONS OF RELEASE

- B**

### CONDITIONS OF RELEASE

- C

### CONDITIONS OF RELEASE

- D

## AMENDMENTS

[illegible]

0065160

CONDITIONS OF RELEASE

BR-113133

\*\*\*\*\*

DRIC U

COPYRIGHT (c)  
1988  
CONTROLLER  
HMSO LONDON

\*\*\*\*\*

DRIC Y

Reports quoted are not necessarily available to members of the public or to commercial organisations.

UNCLASSIFIED/UNLIMITED

ARE TM(UDT)89114  
Accession No 79462

July 1989

A Note On The Gaussian Integration Formula

by

C. Richardson



Accession For	
1000 0001	<input checked="checked" type="checkbox"/>
1000 0002	<input type="checkbox"/>
1000 0003	<input type="checkbox"/>
1000 0004	<input type="checkbox"/>
1000 0005	
1000 0006	
1000 0007	
1000 0008	
1000 0009	
1000 0010	
1000 0011	
1000 0012	
1000 0013	
1000 0014	
1000 0015	
1000 0016	
1000 0017	
1000 0018	
1000 0019	
1000 0020	
1000 0021	
1000 0022	
1000 0023	
1000 0024	
1000 0025	
1000 0026	
1000 0027	
1000 0028	
1000 0029	
1000 0030	
1000 0031	
1000 0032	
1000 0033	
1000 0034	
1000 0035	
1000 0036	
1000 0037	
1000 0038	
1000 0039	
1000 0040	
1000 0041	
1000 0042	
1000 0043	
1000 0044	
1000 0045	
1000 0046	
1000 0047	
1000 0048	
1000 0049	
1000 0050	
1000 0051	
1000 0052	
1000 0053	
1000 0054	
1000 0055	
1000 0056	
1000 0057	
1000 0058	
1000 0059	
1000 0060	
1000 0061	
1000 0062	
1000 0063	
1000 0064	
1000 0065	
1000 0066	
1000 0067	
1000 0068	
1000 0069	
1000 0070	
1000 0071	
1000 0072	
1000 0073	
1000 0074	
1000 0075	
1000 0076	
1000 0077	
1000 0078	
1000 0079	
1000 0080	
1000 0081	
1000 0082	
1000 0083	
1000 0084	
1000 0085	
1000 0086	
1000 0087	
1000 0088	
1000 0089	
1000 0090	
1000 0091	
1000 0092	
1000 0093	
1000 0094	
1000 0095	
1000 0096	
1000 0097	
1000 0098	
1000 0099	
1000 0100	

A-1

• Controller HMSO London, 1989

## CONTENTS

	Page
ABSTRACT ... ..	ii
1. BACKGROUND ... ..	1
2. THEORY ... ..	1
3. FORTRAN PROCEDURE ... ..	5
4. EXAMPLE ... ..	6

## FIGURES

1. Gaussian Integration Of The Function  $\sin(x)$

A Note On The Gaussian Integration Formula

ABSTRACT

The Gaussian integration method is described together with the 2-point implementation for numerical integration. A FORTRAN program is given and the result of a test example.

### BACKGROUND

1. This work was originally completed some years ago and the software, which is incorporated in a Maths Library on the UDT computer facility, has been used extensively. Gauss' formula provides a method for calculating the definite integral of a given function. *Legendre's*

### THEORY

2. If  $P_m(x)$ ,  $P_n(x)$ ,  $x \in \mathbb{R}$ , are polynomials of degree  $m, n$  respectively, with  $m \geq n$ , then polynomials  $q(x)$ ,  $r(x)$  exist such that:

$$P_m(x) = P_n(x)q(x) + r(x) \quad 1.$$

where the degree of  $q(x)$  is  $m-n$  and the degree of  $r(x)$  is  $n-1$  at most. The polynomials  $q(x)$ ,  $r(x)$  are unique and called the quotient and remainder of  $P_m(x)$  with reference to  $P_n(x)$ .

3. Consider:

$$\int_{-1}^{+1} P_{2n+1}(x) dx$$

and let  $P_{n+1}(x)$  be a Legendre polynomial of degree  $n+1$ . Then by equation 1:

$$\int_{-1}^{+1} P_{2n+1}(x) dx = \int_{-1}^{+1} \{P_{n+1}(x)q_n(x) + r_n(x)\} dx \quad 2.$$

But any polynomial of degree  $n$  can be expressed as a linear combination of Legendre polynomials of degree  $k \leq n$ . Thus

$$q_n(x) = \sum_{k=0}^n a_k P_k(x) \quad 3.$$

and therefore, using the orthogonal properties of Legendre functions:

$$\int_{-1}^{+1} P_{n+1}(x)q_n(x) dx = 0 \quad 4.$$

gives:

$$\int_{-1}^{+1} P_{2n+1}(x) dx = \int_{-1}^{+1} r_n(x) dx \quad 5.$$

4. We seek solutions of the form:

$$\begin{aligned} \int_{-1}^{+1} P_{2n+1}(x) dx &= \sum_{k=0}^n a_k P_{2n+1}(x_k) \\ &= \sum_{k=0}^n a_k \{ P_{n+1}(x_k) q_n(x_k) + r_n(x_k) \} \end{aligned} \quad 6.$$

but equation 5 shows the integral is independent of the choice of  $q_n(x)$  and therefore equation 6 is true iff  $x_k$  are the  $n+1$  roots of  $P_{n+1}(x)$ . Let:

$$\omega_{n+1}(x) = \prod_{k=0}^n (x - x_k) \quad 7.$$

then expanding in partial fractions gives:

$$\frac{r_n(x)}{\omega_{n+1}(x)} = \sum_{k=0}^n \frac{r_n(x_k)}{\omega'_{n+1}(x_k)(x - x_k)} \quad 8.$$

Integration of equation 8 gives:

$$\int_{-1}^{+1} r_n(x) dx = \int_{-1}^{+1} \sum_{k=0}^n \frac{\omega_{n+1}(x) r_n(x_k) dx}{\omega'_{n+1}(x_k)(x - x_k)} \quad 9.$$

Therefore:

$$\int_{-1}^{+1} P_{2n+1}(x) dx = \sum_{k=0}^n a_k P_{2n+1}(x_k) \quad 10.$$

where:

$$a_k = \int_{-1}^{+1} \frac{\omega_{n+1}(x) dx}{\omega'_{n+1}(x_k)(x - x_k)} \quad 11.$$

5. If we use the Hermite interpolation formula to represent a function  $f(x)$  by a polynomial of degree  $2n+1$ , the error in the integral formula is given by:

$$e_n = \frac{f^{(2n+2)}(\xi)}{(2n+2)!} \int_{-1}^{+1} \prod_{k=0}^n (x - x_k)^2 dx \quad 12.$$

where  $-1 \leq \xi \leq +1$ .



6. There are  $n+1$  values of  $x_k$ , and hence this method of integration is called the  $n+1$  point method. For the Gaussian 2-point integral formula put  $n=1$  and then  $x_k$   $k=0,1$  are the zeros of  $P_2(x)$ , ie  $x_k$  are given by:

$$(3x^2-1)/2 = 0$$

therefore:

$$x_k = \pm 1/\sqrt{3}$$

and:

$$\omega_{n+1}(x) = (x^2 - 1/3)$$

Hence the weighting factors are:

$$\begin{aligned} a_k &= \int_{-1}^{+1} \frac{x \pm 1/\sqrt{3}}{2x} dx \\ &= 1 \end{aligned}$$

Finally the formula is:

$$\int_{-1}^{+1} f(x) dx = f\left(\frac{1}{\sqrt{3}}\right) + f\left(-\frac{1}{\sqrt{3}}\right) + \frac{f^{(4)}(\xi)}{135} \quad 13.$$

7. In general  $\int_a^b f(x) dx$  can be transformed to the form of equation 13. Furthermore the accuracy can be improved by subdividing the interval  $a,b$ . Let each subinterval have half width  $h$ , where:

$$h = (b-a)/2n$$

Then:

$$\begin{aligned} \int_a^b f(x) dx &= \sum_{k=0}^{n-1} \int_{2kh+a}^{2(k+1)h+a} f(x) dx \\ &= \sum_{k=0}^{n-1} h \int_{-1}^{+1} f(x) dx' \end{aligned}$$

where:

$$x = x'h + h(2k+1) + a$$

and so by equation 13:

$$\int_a^b f(x)dx = h \sum_{k=0}^{n-1} \left\{ f\left(a+h(2k+1)/3\right) + f\left(a+h(2k+1-1)/3\right) \right\} + \frac{h \cdot n \cdot h^4 f^{(4)}(\xi)}{135} \quad 14.$$

8. The error term in equation 14 can be written:

$$e_n = \frac{(b-a)^5 f^{(4)}(\xi)}{2^5 n^4 135}$$

Taking  $2n$  intervals the error will be:

$$\begin{aligned} e_{2n} &= \frac{(b-a)^5 f^{(4)}(\xi)}{2^5 2^4 n^4 135} \\ &= \frac{e_n}{16} \end{aligned}$$

Put:

$$\begin{aligned} \int_a^b f(x)dx &= I_n + e_n \\ &= I_{2n} + e_{2n} \end{aligned}$$

then a closer approximation is given by:

$$\int_a^b f(x)dx = I_{2n} + (I_{2n} - I_n)/15 \quad 15.$$

9. Gauss' integral formula is not suitable to evaluate an integral of a function given in tabular form, but when the function is given as an analytic expression the method is very useful. A FORTRAN procedure is described to compute:

$$I = \int_a^b f(x)dx + \varepsilon$$

where:

$$|\varepsilon| \leq \text{tol}$$

```

REAL FUNCTION GAUSS_INTEGRATION(F,A,B,TOL)

C      Description:
C      Integrates the function F(x) over the range (A,B), where F is a
C      real function (to be provided). The repeated two-point Gauss
C      method is used until successive approximations differ by less
C      than 15*TOL.

C      Author:
C      C Richardson, ARE (Portland)

C      History:
C      Issue 1.0      22 July      1982
C      Mod   1.1      6 August     1985

C      Modifications:
C      1.1      To adopt coding standard

C      Subroutine Arguments:

      REAL          F              ! (R) Function
      REAL          A,B           ! (R) Limits
      REAL          Tol           ! (R) Required accuracy

C      Local Variables:

      REAL          P1,P2,        ! Integration estimates
      REAL          H,            ! Subinterval half width
      REAL          Cd,G
      INTEGER       Nc            ! Number of subintervals

C      Main Entry Point:

      Nc=8                      ! Number of subintervals
      DO WHILE ((Nc .LT. 31).OR.(ABS(P2-P1) .GT. 15.*tol))
      P1=P2                      ! Save last estimate
      H=(B-A)/(2*Nc)             ! Subinterval half-width
      Cd=A-H
      G=H/1.73205081
      P2=0.                      ! Partial sum
      DO M=1,Nc
      Cd=Cd+H+H
      P2=P2+F(Cd-G)
      P2=P2+F(Cd+G)
      ENDDO
      P2=P2*H                    ! Estimated result
      Nc=2*Nc
      ENDDO
      GAUSS_INTEGRATION=P2+(P2-P1)/15.
      RETURN
      END

```

## PROGRAM GAUSS\_INTEGRATION\_TEST

```

C      Description:
C      Test Gaussian integration by computing the integral of Sin(x)
C      over the range (A,B).

C      Author:
C      C Richardson, ARE (Portland)

C      History:
C      Issue 1.0      22 July      1982

C      Local Variables:
      REAL          A,B,          !Integration interval
      P              P            !Workspace

      EXTERNAL F

C      Main Entry Point:
100  FORMAT(A)

      TYPE *, 'The integration interval A,B must be specified'
      A=ASKR('Enter value of A (Degrees)')
      B=ASKR('Enter value of B (Degrees)')

      CALL PLOT_INIT('WINDOW=A4-V')          !Initialise graphics
      CALL CHAR_SIZE(1.5,2.0)
      CALL GRID_LIN_LIN(30.,50.,180.,200.,A,-1.,B,1.,1)
      CALL CURVE(F,A,B,1.)                  !Draw function over A,B
      CALL MOVE((A+B)/2.,-1.)              !Annotate axes
      CALL CHAR_POSN(-7.5,-2.)
      CALL PLOT_TEXT('Angle (Degrees)',15)
      CALL MOVE(A,0.)
      CALL CHAR_POSN(-4.,0.)
      CALL CHAR_ANGLE(90.)
      CALL CHAR_POSN(-3.,0.)
      CALL PLOT_TEXT('Sin(X)',6)
      CALL CHAR_ANGLE(0.)
      P=GAUSS_INTEGRATION(F,A,B,.00001)      !Calculate numerical integral
      P=P/57.29578
      CALL MOVE(A,1.)                      !Annotate graph
      CALL CHAR_POSN(0.,1.5)
      CALL PLOT_TEXT('Numeric Integral=',18)
      CALL PLOT_NUM(P,3,4)
      P=COSD(A)-COSD(B)                    !Calculate analytic integral
      CALL MOVE(A,1.)                      !Annotate graph
      CALL CHAR_POSN(0.,0.5)
      CALL PLOT_TEXT('Analytic Integral=',18)
      CALL PLOT_NUM(P,3,4)
      CALL ORIGIN(0.,0.)
      CALL SCALE(1.,1.)                    !Plot title
      CALL MOVE(30.,20.)
      CALL CHAR_SIZE(1.8,2.4)
      CALL PLOT_TEXT(
      *   'Fig 1. Gaussian Integration Of The Function Sin(X)',51)
      ACCEPT 100,P
      CALL PLOT_FIN
      END

      FUNCTION F(X)                        !A typical function
      F=SIND(X)
      RETURN
      END

```




Fig 1. Gaussian Integration Of The Function Sin(X)